

Hepatic Biotransformation & CYP450 Drug-Drug Interactions (DDIs) (Clinical Treatise 2016)

Por Cristiano Ricardo · Publicado em 27/07/2016

University clinical pharmacology treatise by Prof. Cristiano Ricardo on hepatic biotransformation & cyp450 drug-drug interactions (ddis) (Xenobiotic Metabolism & Molecular Toxicology), featuring mathematical PK/PD modeling, clinical cases, and colorful schematics.

Versão online e eventuais atualizações:

<https://professor.cristianoricardo.com/post/hepatic-biotransformation-cyp450-drug-drug-interactions-ddis-2016-07-27>

Xenobiotic Metabolism & Molecular Toxicology · Clinical Pharmacology & Pharmacotherapy

Enzymatic transformation by the cytochrome P450 superfamily dictates oral first-pass clearance, bioactivation of prodrugs, and severe adverse drug-drug interactions.

Pharmacological Mechanism & Kinetic Schema

Phase I Oxidation

CYP3A4, 2D6, 2C9, 2C19

Introduces -OH, -NH₂

Inhibition ? Toxicity

Phase II Conjugation

UGT (Glucuronidation)

Sulfation & Glutathione

Excretable Polar Metabolite

PXR/CAR Induction

Rifampin, St. John's Wort

Transcriptional boost

Sub-therapeutic failure

Mechanisms of CYP Inhibition: Reversible vs Quasi-Irreversible

Competitive inhibition occurs instantaneously when two co-administered drugs compete for the heme catalytic pocket (e.g., Fluconazole inhibiting CYP2C9, leading to toxic Warfarin accumulation). In contrast, mechanism-based (suicide) inhibition involves covalent binding of reactive metabolic intermediates to the apoprotein (e.g., Clarithromycin inactivating CYP3A4), requiring de novo enzyme synthesis over several days for clinical recovery.

Nuclear Receptor Activation and Transcriptional Enzyme Induction

Potent inducers like Rifampin and Carbamazepine bind to nuclear Pregnane X Receptor (PXR) and Constitutive Androstane Receptor (CAR). Translocation to the nucleus heterodimerizes with Retinoid X Receptor (RXR), massively upregulating CYP3A4, CYP2C9, and P-glycoprotein (ABCB1) transcription, causing therapeutic failure of oral contraceptives and calcineurin inhibitors.

Pharmacogenomic Polymorphisms: The CYP2D6 Spectrum

Genetic copy number variations and single nucleotide polymorphisms categorise patients from Poor Metabolizers (PMs, lacking codeine analgesia due to failure of morphine bio-activation) to Ultra-Rapid Metabolizers (UMs, facing life-threatening opioid toxicity under standard codeine doses).

Prof. Cristiano Ricardo

Clinical Pharmacologist & Biochemist · Specialist in Pharmacokinetics & Hospital Pharmacotherapy · Published on July 27, 2016 at 02:05